



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND
CITIES—UNTABULATED.

AFRICA—*Lourenço Marquez*.—Month of November, 1905. Estimated population, 10,000. Total number of deaths 46, including 8 from tuberculosis.

AUSTRIA-HUNGARY.—*Brünn*.—Month of November, 1905. Estimated population, 95,342. Total number of deaths, 189, including diphtheria 3, enteric fever 2, and 28 from tuberculosis.

BAHAMAS—*Dunmore Town*.—Two weeks ended January 13, 1906. Estimated population, 1,232. One death. No contagious diseases.

Governors Harbor.—Week ended January 13, 1906. Estimated population, 1,500. No deaths and no contagious diseases.

Green Turtle Cay.—Two weeks ended January 11, 1906. Estimated population, 3,314. No deaths and no contagious diseases reported.

Nassau.—Two weeks ended January 13, 1906. Estimated population, 12,650. Number of deaths not reported. No contagious diseases reported.

CANADA—*British Columbia, Victoria*.—Month of December, 1905. Estimated population, 21,000. Total number of deaths, 25, including 2 from tuberculosis.

CHILE—*Antofagasta*.—Month of October, 1905. Estimated population, 24,000. Total number of deaths, 145, including enteric fever 3, plague 7, smallpox 51, and 18 from phthisis pulmonalis.

Month of November, 1905. Total number of deaths, 138, including typhus fever 1, plague 8, smallpox 27, and 13 from phthisis pulmonalis.

FRANCE—*Marseille*.—Month of December, 1905. Estimated population, 491,161. Total number of deaths, 906, including diphtheria 5, enteric fever 15, scarlet fever 1, whooping cough 1, and 101 from tuberculosis.

GERMANY—*Strasburg*.—Month of November, 1905. Estimated population, 165,187. Total number of deaths, 221, including diphtheria 4, measles 4, scarlet fever 1, and 30 from tuberculosis.

GIBRALTAR.—Two weeks ended January 7, 1906. Estimated population, 27,460. Total number of deaths, 10, including 2 from smallpox.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 76 great towns in England and Wales during the week ended January 6, 1906, correspond to an annual rate of 18.3 per 1,000 of the aggregate population, which is estimated at 15,818,360.

London.—One thousand seven hundred and fifty-six deaths were registered during the week, including measles 64, scarlet fever 9, diphtheria 13, whooping cough 33, enteric fever 6, and 10 from diarrhea. The deaths from all causes correspond to an annual rate of 19.4 per 1,000. In Greater London 2,438 deaths were registered. In

the "outer ring" the deaths included 2 from diphtheria, 5 from measles, and 4 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended January 6, 1906, in the 21 principal town districts of Ireland, was 20.6 per 1,000 of the population, which is estimated at 1,093,959. The lowest rate was recorded in Portadown, viz, 5.2, and the highest in Lisburn, viz, 45.5 per 1,000. In Dublin and suburbs 168 deaths were registered, including diphtheria 2, enteric fever 3, whooping cough 2, and 33 from tuberculosis.

Scotland.—The deaths registered in 8 principal towns during the week ended January 6, 1906, correspond to an annual rate of 19.8 per 1,000 of the population, which is estimated at 1,787,788. The lowest rate of mortality was recorded in Paisley, viz, 13.2, and the highest in Dundee, viz, 26.2 per 1,000. The aggregate number of deaths registered from all causes was 680, including measles 33, diphtheria 13, enteric fever 1, scarlet fever 2, and 11 from whooping cough.

ITALY—*Naples*.—Month of November, 1905. Estimated population, 578,371. Total number of deaths, 929, including diphtheria 5, enteric fever 9, measles 1, whooping cough 2, smallpox 1, and 54 from tuberculosis.

JAMAICA—*Kingston*.—Month of December, 1905. Estimated population, 52,475. Total number of deaths, 139, including 8 from phthisis pulmonalis.

MALTA.—Two weeks ended December 30, 1905. Estimated population, 202,134. Total number of deaths, 197, including diphtheria 5, and 1 from enteric fever.

NICARAGUA—*San Juan del Norte*.—Three months ended September 30, 1905. Estimated population, 600. Total number of deaths, 5. No deaths from contagious diseases.

SPAIN—*Cadiz*.—Month of December, 1905. Estimated population, 69,382. Total number of deaths, 180, including diphtheria 2, enteric fever 1, measles 3, whooping cough 1, smallpox 1, and 19 from tuberculosis.